

On the magnetohydrodynamic description of a two-component plasma in the Kerr metric

Ramon Khanna

Landessternwarte, Königstuhl, D-69117 Heidelberg, Germany
rkhanna@lsw.uni-heidelberg.de

5 February 2008

ABSTRACT

The magnetohydrodynamic equations describing an inviscid fully ionized plasma in the vicinity of a rotating black hole are derived from a two-component plasma theory within the framework of the 3+1 split of the Kerr metric. Of central interest is the generalized Ohm's law. In the limit of quasi-neutral plasma it contains no new terms as compared with special relativity. Gravitomagnetic terms appear in Ohm's law only, if the plasma is charged in its rest frame or the MHD-approximation is not applied.

It is argued that a relativistic single-fluid description of a multiple-component plasma is possible only for cold (i.e. intrinsically non-relativistic) components. As seen by local stationary observers, close to the horizon the electron collision time becomes longer than dynamical timescales, i.e. the plasma appears to behave as particles.

Key words: black hole physics – MHD – plasmas – relativity

1 INTRODUCTION

Astrophysics of black hole systems with their magnetized accretion discs, magnetospheres and jets has been a fascinating field of research for some twenty years. There are a number of papers dealing with the general-relativistic description of those systems. The majority of the papers is restricted to ideal MHD of black hole magnetospheres and jets (Blandford & Znajek 1977; Phinney 1983; Bekenstein & Eichler 1985; Camenzind 1986, 1987; Takahashi et al. 1990; Okamoto 1992; Beskin & Par'ev 1993; Fendt 1997). General-relativistic resistive MHD has been discussed by Bekenstein & Oron (1978). Khanna & Camenzind (1994, 1996a,b), Khanna (1997) and Kudoh & Kaburaki (1996) have applied resistive MHD to accretion discs of Kerr black holes. In both ideal and resistive MHD the authors have used the standard Ohm's law.

Generalized Ohm's laws in special-relativistic context have been derived from invariant distribution functions by Ardavan (1976) and by Blackman & Field (1993). Ardavan shows that the simple ideal MHD condition from the standard Ohm's law breaks down in extreme regions of a pulsar magnetosphere. This result suggests that also around a black hole there should be regions in which the standard Ohm's law is merely a poor approximation.

In this paper I derive the full set of MHD equations, including a generalized Ohm's law, for an inviscid, fully ionized plasma from a two-component plasma theory within the framework of the 3+1 split of the Kerr metric. I discuss two major issues: (i) under which circumstances is a relativis-

tic single-fluid description of a multiple-component plasma possible? (ii) Does the gravitomagnetic field play a role in Ohm's law?

The equations and results of Sect. 2 are general and apply also to an electron-positron plasma. The MHD description of Sect. 3 is limited to a relativistically cold electron-ion plasma.

I use a metric with signs $(- + + +)$ and set $c = 1$. Spacetime vectors and tensors are written upright. In 3-dimensional space vectors and tensors are denoted by arrows (e.g. $\vec{v}$ and $\vec{T}$) and covariant derivatives are indicated by vertical bars, e.g. $\nabla_j \beta^i \equiv \beta^i_{|j}$. Superscripts $i, e, '$ indicate the rest frames K^i, K^e and K' , in which a quantity is defined.

2 TWO-COMPONENT PLASMA THEORY IN THE 3+1 SPLIT OF THE KERR METRIC

In this section I will first discuss the relativistic definition of a plasma as the centre-of-mass fluid of its two components and introduce the various rest frames involved. Then I will summarize the 3+1 split of the Kerr metric, in which the equations governing the individual components of the plasma will be written down and finally combined (in Sect. 3) to obtain the equation of motion, the local law of energy conservation and the generalized Ohm's law for the plasma as single fluid.

2.1 Basic concepts, definitions and equations

2.1.1 Ions, electrons and plasma

A fluid of charged particles is described by its stress-energy spacetime tensor, its number-flux 4-vector and its current 4-vector J_x . Thus for interacting ions (in this section ‘ions’ could mean positrons as well) and electrons we have for the hydrodynamic part of the stress-energy spacetime-tensor

$$T_x^{\alpha\beta} = (\rho_x^x + p_x^x)W_x^\alpha W_x^\beta + p_x^x g^{\alpha\beta} + T_{x\text{coll}}^{\alpha\beta}, \quad (1)$$

where $\rho_x^x = n_x m_x (1 + \Pi)$ is the total density of mass-energy (including specific internal energy Π) and p_x^x is the total pressure of the species ‘x’ as measured in the rest frame K^x of the species (x=i,e). W_x is the bulk 4-velocity of the species. The interaction between electrons and ions is described by $T_{x\text{coll}}^{\alpha\beta}$. The number-flux 4-vectors are

$$N_x^\alpha = n_x W_x^\alpha, \quad (2)$$

with the species’ rest frame particle densities n_x .

The plasma is assumed to be a perfect fluid and is defined by the sums of the ion and electron tensors:

$$(\rho_m' + p')W^\alpha W^\beta + p' g^{\alpha\beta} \equiv T^{\alpha\beta} = T_i^{\alpha\beta} + T_e^{\alpha\beta}, \quad (3)$$

$$N^\alpha = N_i^\alpha + N_e^\alpha \quad (4)$$

and

$$J^\alpha = J_i^\alpha + J_e^\alpha. \quad (5)$$

Note that, from Eq. (3), the plasma’s 4-velocity, energy density, momentum density and pressure are uniquely defined as centre-of-mass quantities of ions plus electrons. This means, in particular, that $N^\alpha \neq n_p W^\alpha$, where n_p would be the number density of ‘plasma particles’, since there is no more freedom for such a definition. The consequences of this fact for the possibility of describing the plasma as a single fluid will be discussed below.

Electrodynamic parts of T_x and T are not given here, but will appear as Lorentz force and electric energy in the laws of momentum conservation and energy conservation below.

2.1.2 Rest frames

The plasma rest frame K' is defined as the frame in which the plasma’s centre-of-mass is at rest

$$\vec{v}' = \frac{(\rho_{mi}^i + p_i^i)\vec{v}_i' + (\rho_{me}^e + p_e^e)\vec{v}_e' + \vec{S}_{i\text{coll}}' + \vec{S}_{e\text{coll}}'}{\rho_m' + p'} = \vec{0} \quad (6)$$

(for the definition of momentum densities see below). The idea of a plasma implies that, in the plasma rest frame, both the ion-component (subscript ‘i’) and the electron-component (subscript ‘e’) have non-relativistic *bulk* velocities $\vec{v}_i'$ and $\vec{v}_e'$, i.e. the bulk Lorentz factors $\gamma_i' = 1$ and $\gamma_e' = 1$ (this does not mean that the plasma components may not be relativistically hot). In this sense the rest frames of electrons K^e , ions K^i and plasma K' are identical and quantities defined in K^e and K^i will be the same in K' to high accuracy.

The total densities of mass-energy as measured in K' are

$$\epsilon_i' = (\rho_{mi}^i + p_i^i v_i'^2) \gamma_i'^2 + \epsilon_{i\text{coll}}' \approx \rho_{mi}^i$$

$$\begin{aligned} \epsilon_e' &= (\rho_{me}^e + p_e^e v_e'^2) \gamma_e'^2 + \epsilon_{e\text{coll}}' \approx \rho_{me}^e \\ \epsilon' &= \rho_m' = \epsilon_i' + \epsilon_e' \approx \rho_{mi}^i + \rho_{me}^e, \end{aligned} \quad (7)$$

where the approximations are due to $v_x'^2/c^2 \ll 1$ and the energy exchange in collisions being small compared to the rest mass of the particles. The latter is true for a plasma in equilibrium, i.e. behaving as one fluid. In this case collisions between electrons and ions are nearly elastic and also the energy flux $\vec{S}_{x\text{coll}}'$ is negligible. The pressures $p_i' \simeq p_i^i$ and $p_e' \simeq p_e^e$ add up to the total pressure

$$p' = p_i' + p_e'. \quad (8)$$

The charge densities are

$$\rho_{ci}' \simeq \rho_{ci}^i = Z n_i \quad \rho_{ce}' \simeq \rho_{ce}^e = -e n_e \quad \rho_c' = \rho_{ci}' + \rho_{ce}'. \quad (9)$$

2.1.3 The 3+1 split of the Kerr metric

In the 3+1 split of the Kerr metric (see Thorne et al. 1986), spacetime is split into a family of three dimensional differentially rotating hypersurfaces of constant time with internal curvature. Those hypersurfaces can be mentally collapsed into a single 3-dimensional ‘absolute space’ in which time is globally described by the Boyer-Lindquist coordinate t . The line element of the Kerr metric in 3+1 notation is given by

$$\begin{aligned} ds^2 &= g_{\alpha\beta} dx^\alpha dx^\beta \\ &= -\alpha_g^2 dt^2 + h_{jk}(dx^j + \beta^j dt)(dx^k + \beta^k dt), \end{aligned} \quad (10)$$

where the lapse function is identified with the gravitational redshift

$$\alpha_g \equiv \left(\frac{d\tau}{dt} \right)_{\text{FIDO}} = \frac{\rho}{\Sigma} \sqrt{\Delta}, \quad (11)$$

τ being the local proper time as measured by FIDOs (see below). The shift functions are the components of the gravitomagnetic potential $\vec{\beta}$

$$\beta^r = \beta^\theta = 0, \quad \beta^\phi \equiv -\omega = -\left(\frac{d\phi}{dt} \right)_{\text{FIDO}} = -\frac{2aMr}{\Sigma^2}, \quad (12)$$

and the components of the 3-metric $\vec{h}$ are

$$h_{rr} = \frac{\rho^2}{\Delta}, \quad h_{\theta\theta} = \rho^2, \quad h_{\phi\phi} = \tilde{\omega}^2, \quad h_{jk} = 0; (j \neq k). \quad (13)$$

Note that $h_{jk} = g_{jk}$ but $h^{jk} = g^{jk} + \beta^j \beta^k / \alpha_g^2$. The metric functions appearing here are defined as

$$\begin{aligned} \Delta &\equiv r^2 - 2Mr + a^2, \quad \rho^2 \equiv r^2 + a^2 \cos^2 \theta, \\ \Sigma^2 &\equiv (r^2 + a^2)^2 - a^2 \Delta \sin^2 \theta, \quad \tilde{\omega} \equiv (\Sigma/\rho) \sin \theta. \end{aligned} \quad (14)$$

The parameters of the black hole are its mass M and angular momentum J , which define the Kerr parameter $a \equiv J/M$.

2.1.4 Quantities and equations in the FIDO frame

Fiducial observers (FIDOs) who live in ‘absolute Kerr-space’ are stationary and locally non-rotating zero angular momentum observers (ZAMOs). As seen in a global Boyer-Lindquist coordinate system, ZAMOs are in shearing circular motion around the Kerr black hole, described by their 4-velocity

$$U = \alpha_g^{-1} (\vec{e}_t - \vec{\beta}). \quad (15)$$

The quantities and equations describing physics in the locally flat laboratories of the FIDOs are obtained from spacetime vectors, tensors and tensorial equations by projection along $\mathbf{U}$ and orthogonal to $\mathbf{U}$ with the projection operator

$$h_{\alpha\beta} = g_{\alpha\beta} + U_\alpha U_\beta . \quad (16)$$

The relevant plasma quantities in the context of this paper are, for ions and electrons respectively, the bulk velocities $\vec{v}_i$, $\vec{v}_e$ and Lorentz factors γ_i , γ_e , the energy densities

$$\epsilon_x = (\rho_{mx}^x + p_x^x \gamma_x^2) \gamma_x^2 + \epsilon_{x \text{ coll}} \approx (\rho_{mx}^x + p_x^x \gamma_x^2) \gamma_x^2 \quad (17)$$

and the momentum densities

$$\vec{S}_x = (\rho_{mx}^x + p_x^x) \gamma_x^2 \vec{v}_x + \vec{S}_{x \text{ coll}} \approx (\rho_{mx}^x + p_x^x) \gamma_x^2 \vec{v}_x . \quad (18)$$

The stress-energy tensor for an interacting fluid in ‘absolute space’ is given by

$$\vec{T}_x = (\rho_{mx}^x + p_x^x) \gamma_x^2 \vec{v}_x \otimes \vec{v}_x + p_x^x \vec{h} + \vec{T}_{x \text{ coll}} . \quad (19)$$

The plasma can be treated as a single fluid, if collisions between ions and electrons involve energies and energy fluxes, much smaller than the rest mass of the particles. Then the collision terms can be neglected in Eqs. (17) and (18). In $\vec{T}_x$ the collision term has to be kept, however, as it describes the momentum flux, which couples ions and electrons. Note that ϵ_x , $\vec{S}_x$ and $\vec{T}_x$ result from splitting $\mathbf{T}_x$.

The charge densities and current densities of the plasma components are, respectively,

$$\rho_{ci} = \rho_{ci}^i \gamma_i = \frac{\rho_{ci}^i}{\gamma(1 - \vec{v} \cdot \vec{v}_i)} \quad \rho_{ce} = \rho_{ce}^e \gamma_e = \frac{\rho_{ce}^e}{\gamma(1 - \vec{v} \cdot \vec{v}_e)} \quad (20)$$

$$\vec{j}_i = \rho_{ci} \vec{v}_i \quad \vec{j}_e = \rho_{ce} \vec{v}_e . \quad (21)$$

Equations (20) are derived in appendix A.

In the context of a two-component theory, each plasma component (subscript $x=i,e$) has to obey the continuity equation for the density of rest-mass (see appendix B)

$$\left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) (n_x m_x \gamma_x) + \vec{\nabla} \cdot (\alpha_g n_x m_x \gamma_x \vec{v}_x) = 0 \quad (22)$$

and the law of momentum conservation (Thorne et al. 1986; Thorne & Macdonald 1982)

$$\frac{1}{\alpha_g} \left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) \vec{S}_{x \text{ id}} = \epsilon_{x \text{ id}} \vec{g} + \vec{H} \cdot \vec{S}_{x \text{ id}} - \frac{1}{\alpha_g} \vec{\nabla} \cdot (\alpha_g \vec{T}_{x \text{ id}}) + \rho_{cx} \vec{E} + \vec{j}_x \times \vec{B} + \vec{k}_x , \quad (23)$$

where

$$\begin{aligned} \vec{k}_x &= \epsilon_{x \text{ coll}} \vec{g} + \vec{H} \cdot \vec{S}_{x \text{ coll}} - \frac{1}{\alpha_g} \vec{\nabla} \cdot (\alpha_g \vec{T}_{x \text{ coll}}) \\ &- \frac{1}{\alpha_g} \left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) \vec{S}_{x \text{ coll}} \approx - \frac{1}{\alpha_g} \vec{\nabla} \cdot (\alpha_g \vec{T}_{x \text{ coll}}) \end{aligned} \quad (24)$$

is an interaction force density due to collisions between ions and electrons. Note that, on physical grounds, $\vec{k}_i = -\vec{k}_e$. The subscript ‘id’ refers to the ideal (inviscid) part of the quantity. Equation (23) contains all the terms known from flat space plus an accelerating force along $\vec{\beta}$ on the l.h.s. plus, on the r.h.s., a coupling term between the gravitomagnetic tensor field

$$\vec{H} \equiv \alpha_g^{-1} \vec{\nabla} \vec{\beta}, \quad H_{ij} = \alpha_g^{-1} \beta_{j|i} \quad (25)$$

and $\vec{S}_x$, which acts as gravitomagnetic force similar to the electromagnetic Lorentz force. The gravitational acceleration is related to the gravitational redshift by $\vec{g} = -\vec{\nabla} \ln \alpha_g$.

The magnetohydrodynamic law of energy conservation for each individual fluid is (Thorne et al. 1986; Thorne & Macdonald 1982)

$$\frac{1}{\alpha_g} \left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) \epsilon_x = -\vec{\nabla} \cdot \vec{S}_x + 2\vec{g} \cdot \vec{S}_x + \vec{H} : \vec{T}_x + \vec{j}_x \cdot \vec{E} . \quad (26)$$

The term $\vec{H} : \vec{T}_x \equiv H_{ij} T_x^{ij}$ can be interpreted as the rate per unit volume at which the gravitomagnetic field does work on matter.

Furthermore the electrodynamics of the plasma is described by Maxwell’s equations and the law of charge conservation (Thorne et al. 1986)

$$\left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) \rho_{cx} + \vec{\nabla} \cdot (\alpha_g \vec{j}_x) = 0 . \quad (27)$$

3 ONE-FLUID DESCRIPTION OF A COLD ELECTRON-ION PLASMA

In this section the two sets of equations for ions (not positrons) and electrons will be combined to yield the equations of MHD, which should no longer contain quantities of the individual components. It is argued that this is possible only for a cold plasma. ‘Cold’ means intrinsically non-relativistic, i.e. $p_x^x \ll \rho_{mx}^x c^2$, $\rho_{mx}^x \approx n_x m_x$ (specific internal energy density negligible compared to rest mass density) and $\rho_m^i \approx n_i m_i + n_e m_e$. This limit has to be taken of Eqs. (1) and (3), or, equivalently, in Eqs. (6), (7), (17) – (19).

3.1 Particle conservation

Describing a plasma as a single fluid means that one has to define ‘plasma particles’ (subscript ‘p’), which have to fulfil particle conservation

$$\left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) (n_p \gamma_p) + \vec{\nabla} \cdot (\alpha_g n_p \gamma_p \vec{v}_p) = 0 , \quad (28)$$

or $N_{;\alpha}^\alpha \equiv (n_p W_p^\alpha)_{;\alpha} = 0$. As was mentioned in Sect. 2.1.1, identifying $\vec{v}_p$ with the centre-of-mass velocity $\vec{v}$ (or W_p^α with W^α) would be in conflict with Def. (3). This means that, without any further assumptions, ‘plasma particle’ conservation can not be formulated in terms of $\vec{v}$. Adding the continuity equations (22) ($x=i,e$) of the two components rather than the equations of particle conservation yields

$$\begin{aligned} &\left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) (n_i m_i \gamma_i + n_e m_e \gamma_e) \\ &+ \vec{\nabla} \cdot (\alpha_g n_i m_i \gamma_i \vec{v}_i + \alpha_g n_e m_e \gamma_e \vec{v}_e) = 0 . \end{aligned} \quad (29)$$

We want to eliminate the γ_i , γ_e , $\vec{v}_i$ and $\vec{v}_e$ in accordance with Definition (3) and derive a set of equations that contains only quantities of a one-fluid description. $T^{\alpha\beta} = T_i^{\alpha\beta} + T_e^{\alpha\beta}$ splits into

$$\epsilon = \epsilon_i + \epsilon_e \quad (30)$$

and

$$\vec{S} = \vec{S}_i + \vec{S}_e . \quad (31)$$

Definition (3) implies that

$$\epsilon \equiv (\rho'_m + p' v^2) \gamma^2 \approx \rho'_m \gamma^2, \quad (32)$$

$$\vec{S} \equiv (\rho'_m + p') \gamma^2 \vec{v} \approx \rho'_m \gamma^2 \vec{v} \quad (33)$$

like obviously a FIDO would define energy density and momentum density of a plasma with velocity $\vec{v}$ and Lorentz factor γ anyway. In the cold plasma limit and with $\gamma_e \approx \gamma_i \approx \gamma$, the last five equations, together with Eqs. (17) and (18), yield

$$\left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) (\rho'_m \gamma) + \vec{\nabla} \cdot (\alpha_g \rho'_m \gamma \vec{v}) = 0. \quad (34)$$

It is important to note that this continuity equation for the plasma, expressed in terms of centre-of-mass quantities, could only be derived in the cold plasma limit. This suggests (though a strict proof has not been given) that a hot multiple-component plasma may not be described as a single fluid at all. This has far reaching consequences for modelling hot astrophysical plasmas in jets and in hot transsonic accretion flows onto black holes.

Note that $\gamma_e \approx \gamma_i$ is not really a new assumption, but is a consequence of $\gamma'_e = \gamma'_i = 1$. Furthermore, with the cold plasma assumption and $n_i m_i \gg n_e m_e$, Eqs. (30), (32) and Eqs. (31), (33) imply

$$\gamma^2 \approx \gamma_i^2 \text{ and } \gamma^2 \vec{v} \approx \gamma_i^2 \vec{v}_i. \quad (35)$$

3.2 Equation of motion

In the cold plasma limit $\vec{T}_{xid}$ can be eliminated from Eq. (23), using Eq. (22), to yield the equation of motion for each plasma component:

$$\begin{aligned} \rho_{mx}^x \gamma_x \left[\frac{1}{\alpha_g} \frac{\partial}{\partial t} + \left(\vec{v}_x - \frac{\vec{\beta}}{\alpha_g} \right) \cdot \vec{\nabla} \right] (\gamma_x \vec{v}_x) &= \rho_{mx}^x \gamma_x^2 \vec{g} \\ + \rho_{mx}^x \gamma_x^2 \vec{H} \cdot \vec{v}_x - \frac{1}{\alpha_g} \vec{\nabla} \cdot (\alpha_g p_x^x) &+ \rho_{cx} \vec{E} + \vec{j}_x \times \vec{B} + \vec{k}_x \end{aligned} \quad (36)$$

The equation of motion for an inviscid electron-ion plasma in one-fluid description follows by adding Eqs. (36) ($x=i,e$):

$$\begin{aligned} \rho'_m \gamma \left[\frac{1}{\alpha_g} \frac{\partial}{\partial t} + \left(\vec{v} - \frac{\vec{\beta}}{\alpha_g} \right) \cdot \vec{\nabla} \right] (\gamma \vec{v}) &= \rho'_m \gamma^2 \vec{g} \\ + \rho'_m \gamma^2 \vec{H} \cdot \vec{v} - \frac{1}{\alpha_g} \vec{\nabla} \cdot (\alpha_g p') &+ \rho_c \vec{E} + \vec{j} \times \vec{B}. \end{aligned} \quad (37)$$

It does not come as a surprise that this equation looks just like Eq. (36). This is, however, not trivial and includes definitions and approximations that require thorough discussion, in particular in the relativistic context. This will be the topic of the rest of this section. Note that Eq. (37) could be derived straightforwardly by adding up the momentum equations (see Sect. 3.2.1 below) and eliminating $\vec{T} = \vec{T}_i + \vec{T}_e$ with the aid of Eq. (34). This way, however, the physics behind the assumptions that have to be made (or that have been made already) would remain obscure.

As has been elaborated in the last section, energy densities and momentum densities add according to Eq. (3). Adding the current density 4-vectors of each plasma component and splitting into charge density and current density yields

$$\rho_c \equiv \rho_{ci} + \rho_{ce} = Z n_i \gamma_i - e n_e \gamma_e, \quad (38)$$

$$\vec{j} \equiv \vec{j}_i + \vec{j}_e = Z n_i \gamma_i \vec{v}_i - e n_e \gamma_e \vec{v}_e. \quad (39)$$

The expressions for ρ_c and $\vec{j}$ involve Eqs. (9), (20) and (21). Note that the collision force densities have cancelled in Eq. (37).

Besides using the definitions above, there is a fundamental assumption that has been made to arrive at the l.h.s. of Eq. (37). It has been assumed that the two plasma components are coupled sufficiently strong that their bulk accelerations

$$\frac{d(\gamma_x \vec{v}_x)}{d\tau_x} \equiv \left[\frac{\gamma_x}{\alpha_g} \frac{\partial}{\partial t} + \gamma_x \left(\vec{v}_x - \frac{\vec{\beta}}{\alpha_g} \right) \cdot \vec{\nabla} \right] (\gamma_x \vec{v}_x) \quad (40)$$

are nearly the same (standard assumption for the applicability of MHD), i.e.

$$\left| \frac{d(\gamma_i \vec{v}_i)}{d\tau_i} - \frac{d(\gamma_e \vec{v}_e)}{d\tau_e} \right| \ll \left| \frac{d(\gamma_i \vec{v}_i)}{d\tau_i} \right|, \quad (41)$$

where τ_x is the proper time in the frame K^x . The application of this assumption (see Schlüter 1950; Kippenhahn & Möllenhoff 1975 for the non-relativistic equivalent) in the derivation of Eq. (37) is elaborated in appendix (C). In Kerr metric we have to go even a step further for consistency. The acceleration due to the gravitomagnetic force is of the same order as the acceleration along $\vec{\beta}$. In fact, if both gravitomagnetic terms of Eq. (36) are added together their axisymmetric toroidal components cancel. If inequality (41) were applied alone in the derivation of the generalized Ohm's law (see below and appendix E), such an axisymmetric toroidal term would remain as an artefact dynamo term. We therefore also have to assume that the coupling between ions and electrons synchronizes the gravitomagnetic accelerations, i.e.

$$\left| \vec{H} \cdot (\gamma_i^2 \vec{v}_i) - \vec{H} \cdot (\gamma_e^2 \vec{v}_e) \right| \ll \left| \vec{H} \cdot (\gamma_i^2 \vec{v}_i) \right|. \quad (42)$$

This assumption has no effect on the equation of motion, but is crucial in the derivation of Ohm's law.

3.2.1 Exact law of momentum conservation for a two-component plasma

As was mentioned above, Eqs. (23) (with $x=i,e$) can be added to give the exact law of momentum conservation for a two-component plasma

$$\begin{aligned} \frac{1}{\alpha_g} \left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) \vec{S} &= \epsilon \vec{g} + \vec{H} \cdot \vec{S} - \frac{1}{\alpha_g} \vec{\nabla} \cdot (\alpha_g \vec{T}) \\ + \rho_c \vec{E} + \vec{j} \times \vec{B}. \end{aligned} \quad (43)$$

Here $\epsilon, \vec{S}$ may be the full expressions for a hot plasma. The stress-energy space-tensor of the plasma is

$$\vec{T} \equiv (\rho'_m + p') \gamma^2 \vec{v} \otimes \vec{v} + p' \vec{h} \approx \rho'_m \gamma^2 \vec{v} \otimes \vec{v} + p' \vec{h} \quad (44)$$

in consistence with Def. (3). The fact that, in the cold plasma limit, Eqs. (43) and (34) can be combined to yield Eq. (37), without explicit use of inequality (41), implies that the collision terms have to implicitly guarantee synchronized bulk acceleration of the components for Def. (3) to be self-consistent in the first place. In the cold plasma limit, Eq. (43) can be used alternatively to Eq. (37). This is particularly appropriate for numerical treatments of MHD, which requires the solutions of a conservative system of differential equations.

3.3 The law of energy conservation

It is straightforward to show that the laws of energy conservation for the individual plasma components add linearly to give

$$\frac{1}{\alpha_g} \left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) \epsilon = -\vec{\nabla} \cdot \vec{S} + 2\vec{g} \cdot \vec{S} + \vec{H} : \vec{T} + \vec{j} \cdot \vec{E}. \quad (45)$$

Like Eq. (43) this equation is exact, but it makes sense as part of a complete set of equations only with the single-cold-fluid assumptions discussed above.

3.4 The generalized Ohm's law for an electron-ion plasma

Multiplying the equation of motion for the ions by $Zm_e\gamma_e^2$ and the equation of motion for the electrons by $m_i\gamma_i^2$ and subtracting the latter from the former yields

$$\begin{aligned} m_i m_e \left[Z\gamma_e^2 n_i \frac{d(\gamma_i \vec{v}_i)}{d\tau_i} - \gamma_i^2 n_e \frac{d(\gamma_e \vec{v}_e)}{d\tau_e} \right] \\ = m_i m_e \gamma_i^2 \gamma_e^2 \left[(Zn_i - n_e) \vec{g} + \vec{H} \cdot (Zn_i \vec{v}_i - n_e \vec{v}_e) \right] \\ - \frac{Zm_e \gamma_e^2}{\alpha_g} \vec{\nabla}(\alpha_g p_i^i) + \frac{m_i \gamma_i^2}{\alpha_g} \vec{\nabla}(\alpha_g p_e^e) \\ + (Zm_e \gamma_e^2 \rho_{ci} - m_i \gamma_i^2 \rho_{ce}) \vec{E} + (Zm_e \gamma_e^2 \vec{j}_i - m_i \gamma_i^2 \vec{j}_e) \times \vec{B} \\ + Zm_e \gamma_e^2 \vec{k}_i - m_i \gamma_i^2 \vec{k}_e. \end{aligned} \quad (46)$$

This equation can be simplified by dropping terms of the order m_e/m_i as compared to their companion terms: With $O(\gamma_i^2) = O(\gamma_e^2)$ and the assumptions $p_i^i/p_e^e \ll m_i/m_e$, $\rho_{ci}/\rho_{ce} \ll m_i/m_e$, the terms with p_i^i , ρ_{ci} and $\vec{k}_i$ can be neglected. The l.h.s. of the equation and the gravitomagnetic term are approximated according to inequalities (41) and (42), respectively. Finally the term $(\#) \times \vec{B}$ can be manipulated as elaborated in appendix (D) to arrive at

$$\begin{aligned} \frac{m_e(Zn_i \gamma_e^2 - n_e \gamma_i^2)}{en_e \gamma_e \gamma_i^2} \left(\frac{d(\gamma \vec{v})}{d\tau_p} - \vec{H} \cdot (\gamma^2 \vec{v}) \right) \\ \approx \vec{E} + \frac{Zn_i \gamma_i}{n_e \gamma_e} \vec{v} \times \vec{B} - \frac{\vec{j} \times \vec{B}}{en_e \gamma_e} + \frac{\vec{\nabla}(\alpha_g p_e^e)}{en_e \gamma_e \alpha_g} \\ + \frac{m_e \gamma_e}{en_e} (Zn_i - n_e) \vec{g} - \frac{1}{en_e \gamma_e} \vec{k}_e, \end{aligned} \quad (47)$$

where τ_p is the proper time in the plasma frame K' .

In analogy to the classical two-component theory and consistent with the special relativistic treatment of Blackman & Field (1993) one can make the ansatz (motivated by the fact that $\vec{k}_e$ is due to the average momentum transfer between electrons and ions)

$$\vec{k}_e = n_e \nu_c m_e (\gamma_i \vec{v}_i - \gamma_e \vec{v}_e). \quad (48)$$

The effective electron collision frequency ν_c (electron-ion, electron-electron, or electron-wave collisions as measured in the plasma rest frame) may be anisotropic. Classically anisotropy of ν_c is due to the reduced mobility of charged particles perpendicular to magnetic fields. It can be expected that, as gravity, the gravitomagnetic force gives rise to drift motions and therefore ν_c might depend on the direction relative to the gravitomagnetic field as well. Additionally, the gravitomagnetic force causes gyration of particles around the gravitomagnetic field $\vec{H} = \alpha_g^{-1} \vec{\nabla} \times \vec{B}$

with a gyrofrequency as measured in the FIDO frame, $\sim |\vec{H}|$. If particles gyrate freely between collisions, i.e. if $|\vec{H}|/\nu_c \sim (\omega/\alpha_g)/(\nu_c/\gamma) \gg 1$ the collision rate should become anisotropic. A rough estimate of this ratio will be given below. Further discussion of an anisotropic collision rate is beyond the scope of this paper.

Re-expressing $\gamma_i \vec{v}_i$ and $\gamma_e \vec{v}_e$ in Eq. (48) with the aid of Eqs. (D4) and (D3) yields

$$\vec{k}_e = \nu_c m_e (\vec{j} - \rho_c' \gamma \vec{v})/e. \quad (49)$$

Inserting this into Eq. (47), one arrives at the *Generalized Ohm's law*

$$\begin{aligned} \frac{\vec{j}}{\sigma \gamma_e} \approx \vec{E} + \frac{Zn_i \gamma_i}{n_e \gamma_e} \vec{v} \times \vec{B} - \frac{\vec{j} \times \vec{B}}{en_e \gamma_e} + \frac{\vec{\nabla}(\alpha_g p_e^e)}{en_e \gamma_e \alpha_g} \\ + \frac{4\pi \gamma_e}{\omega_{pe}^2} \rho_c' \vec{g} + \frac{\rho_c' \gamma \vec{v}}{\sigma \gamma_e} \\ - \frac{4\pi e (Zn_i \gamma_e^2 - n_e \gamma_i^2)}{\omega_{pe}^2 \gamma_e \gamma_i^2} \left(\frac{d(\gamma \vec{v})}{d\tau_p} - \vec{H} \cdot (\gamma^2 \vec{v}) \right), \end{aligned} \quad (50)$$

where the conductivity

$$\sigma = \frac{e^2 n_e}{m_e \nu_c} \equiv \frac{\omega_{pe}^2}{4\pi \nu_c} \quad (51)$$

and ω_{pe} is the electron plasma frequency. The appearance of the convection current $\rho_c' \gamma \vec{v}$ is compulsory to fulfil the transformation law of the current 4-vector and thereby confirms the correctness of the ansatz made for $\vec{k}_e$ in Eq. (48).

The generalized Ohm's law derived here contains no 'current acceleration' term (see appendix E). This is a consequence of inequalities (41), which has to be applied here in order to be consistent with Eq. (37), i.e. to stay in the MHD regime. The current acceleration term is, however, important only on time-scales in the plasma frame of the order of the collision time of the electrons, e.g. switch-on processes. There is, however, something like an 'acceleration current', which, after multiplication with $\sigma \gamma_e$ and for $\gamma_e \approx \gamma_i$, can be written as

$$\vec{j}_{acc} \approx -\frac{\rho_c'}{\nu_c} \frac{d(\gamma \vec{v})}{d\tau_p}. \quad (52)$$

A similar term can be found in Ardavan (1976) for the special-relativistic case and in Kippenhahn & Möllenhoff (1975) for the non-relativistic case. Both the 'acceleration current' and the 'gravitomagnetic current'

$$\vec{j}_{gm} \approx \frac{\rho_c'}{\nu_c} \vec{H} \cdot (\gamma^2 \vec{v}) \quad (53)$$

appear only in a charged plasma.

3.4.1 The quasi-neutral limit

In the limit of quasi-neutral plasma ($Zn_i \approx n_e$) and $\gamma_e \approx \gamma_i \approx \gamma$ Eq. (50) reduces to

$$\vec{j} \approx \sigma \gamma (\vec{E} + \vec{v} \times \vec{B}) - \frac{\sigma}{en_e} (\vec{j} \times \vec{B}) + \frac{\sigma}{en_e \alpha_g} \vec{\nabla}(\alpha_g p_e^e) \quad (54)$$

which contains all the terms, familiar from the non-relativistic generalized Ohm's law. The relative importance of the Hall-term can be estimated by writing

$$\frac{\sigma}{en_e c} |\vec{j} \times \vec{B}| = \frac{\omega_{ce}}{(\nu_c/\gamma)} \left| \vec{j} \times \frac{\vec{B}}{|\vec{B}|} \right| \leq \frac{\omega_{ce}}{(\nu_c/\gamma)} |\vec{j}|. \quad (55)$$

This shows that the Hall-term gives significant modification of the current, if the electron gyrofrequency measured in the FIDO frame, $\omega_{ce} = e|\vec{B}|/\gamma m_e c$, is of the order or higher than the electron collision rate. According to Spitzer (1962) in a non-magnetized plasma

$$\nu_c \approx 1.5 \cdot 10^{-6} \left(\frac{n_e}{\text{cm}^{-3}} \right) \left(\frac{T}{\text{eV}} \right)^{-3/2} \ln \Lambda \text{ s}^{-1}, \quad (56)$$

where $\ln \Lambda$ is the Coulomb logarithm. For polytropic ($\Gamma = 5/3$) spherical accretion onto a non-rotating black hole (the solutions in Kerr metric will be similar) we have (e.g. Shapiro & Teukolski 1983)

$$n_e \approx 2 \cdot 10^{15} M_1^{-1} \frac{\dot{m}}{0.01} \left(\frac{r}{r_g} \right)^{-3/2} \text{ cm}^{-3} \quad (57)$$

$$T \approx 2 \cdot 10^{12} \left(\frac{r}{r_g} \right)^{-1} \text{ K}, \quad (58)$$

with $r_g = GM/c^2$ and $M_1 \equiv M/10M_\odot$. The dimensionless accretion rate $\dot{m}$ is measured in units of the Eddington rate $\dot{M}_{\text{Edd}} \equiv L_{\text{Edd}}/c^2$. Allowing for cooling, the temperature of a realistic accretion flow will rather be $\sim 10^{10} \text{ K}$. Thus we have

$$\nu_c \approx 3 \cdot 10^1 M_1^{-1} \frac{\dot{m}}{0.01} \frac{\ln \Lambda}{10} \text{ s}^{-1} \quad (59)$$

and

$$\frac{\omega_{ce}}{(\nu_c/\gamma)} \approx 6 \cdot 10^2 M_1 \left(\frac{|\vec{B}|}{\text{mGauss}} \right) \left(\frac{\dot{m}}{0.01} \right)^{-1} \left(\frac{\ln \Lambda}{10} \right)^{-1}. \quad (60)$$

Obviously, at those high temperatures and low collision rates, the Hall-term would dominate the current even at low field strength. Note, however, that in such a situation cross-field driven plasma instabilities are likely to occur, resulting in an increased anomalous collision frequency.

As was mentioned above, also gravitomagnetic drifts might cause anisotropic collision rates. The gravitomagnetic gyrofrequency is given by

$$|\vec{H}| \sim \omega = \frac{c}{2r_g} \left(\frac{2r_g \omega}{c} \right) = 10^4 M_1^{-1} \left(\frac{2r_g \omega}{c} \right) \text{ s}^{-1}. \quad (61)$$

The ratio of gravitomagnetic gyrofrequency to Spitzer's collision rate is, as measured in FIDO frames,

$$\frac{(\omega/\alpha_g)}{(\nu_c/\gamma)} \sim 3 \cdot 10^2 \frac{\gamma}{\alpha_g} \left(\frac{2r_g \omega}{c} \right) \left(\frac{\dot{m}}{0.01} \right)^{-1} \left(\frac{\ln \Lambda}{10} \right)^{-1}. \quad (62)$$

This result hints to a problem: ω is the inverse of a dynamical timescale. By applying the MHD-approximation (Eqs. [41] and [42]), which has led to neglecting the current acceleration term, we have implicitly made the assumption that our timescales of interest are much larger than the electron collision time. At very high temperatures Spitzer's collision rate is therefore too low to facilitate single-fluid behaviour of the plasma. Thus the gravitomagnetic field should introduce anisotropy of ν_c and of the particle distribution function. An anisotropic distribution function could trigger plasma instabilities with anomalous collision rates, which should make $(\omega/\alpha_g)/(\nu_c/\gamma) < 1$. At the horizon the ratio of locally measured gravitomagnetic gyro- to collision frequency blows up as $\gamma/\alpha_g \rightarrow \infty$, which means that the MHD description breaks down for FIDOs, i.e. the plasma appears to behave as test particles.

3.5 Summary of equations

As was argued above, a closed set of equations of MHD can only be formulated in the cold plasma limit with $\gamma_e \approx \gamma_i \approx \gamma$. The complete set of equations for a electron-ion plasma are: The continuity equation

$$\left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) (\rho'_m \gamma) + \vec{\nabla} \cdot (\alpha_g \rho'_m \gamma \vec{v}) = 0, \quad (63)$$

with $\rho'_m \approx n_i m_i$.

The local law of momentum conservation

$$\begin{aligned} \frac{1}{\alpha_g} \left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) \vec{S} &= \epsilon \vec{g} + \vec{H} \cdot \vec{S} - \frac{1}{\alpha_g} \vec{\nabla} \cdot (\alpha_g \vec{T}) \\ &+ \rho_c \vec{E} + \vec{j} \times \vec{B}, \end{aligned} \quad (64)$$

with $\epsilon \approx \rho'_m \gamma^2$, $\vec{S} \approx \rho'_m \gamma^2 \vec{v}$ and $\vec{T} \approx \rho'_m \gamma^2 \vec{v} \otimes \vec{v} + p' \vec{h}$. The generalized Ohm's law

$$\begin{aligned} \frac{\vec{j}}{\sigma \gamma} &\approx \vec{E} + \frac{Z n_i}{n_e} \vec{v} \times \vec{B} - \frac{\vec{j} \times \vec{B}}{e n_e \gamma} + \frac{\vec{\nabla}(\alpha_g p_e^e)}{e n_e \gamma \alpha_g} \\ &+ \frac{4\pi \gamma}{\omega_{pe}^2} \rho'_c \vec{g} + \frac{\rho'_c \gamma \vec{v}}{\sigma \gamma} \\ &- \frac{4\pi \rho'_c}{\omega_{pe}^2 \gamma} \left(\frac{d(\gamma \vec{v})}{d\tau_p} - \vec{H} \cdot (\gamma^2 \vec{v}) \right). \end{aligned} \quad (65)$$

Note that $n_e \approx Z \rho'_m / m_i - \rho'_c / e$ and ρ'_c has to be calculated from Eq. (A1). The electron pressure follows, e.g., with the assumption that electrons and ions have the same temperature. Then $p_e^e \approx Z p' / (Z + 1)$.

The local law of energy conservation

$$\frac{1}{\alpha_g} \left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) \epsilon = -\vec{\nabla} \cdot \vec{S} + 2\vec{g} \cdot \vec{S} + \vec{H} \cdot \vec{T} + \vec{j} \cdot \vec{E}. \quad (66)$$

The law of charge conservation

$$\left(\frac{\partial}{\partial t} - \vec{\beta} \cdot \vec{\nabla} \right) \rho_c + \vec{\nabla} \cdot (\alpha_g \vec{j}) = 0. \quad (67)$$

The set of equations is completed by Maxwell's equations (Thorne et al. 1986), from which ρ_c and $\vec{j}$ can be computed without having to solve for $\vec{v}_x$ in the two-fluid theory:

$$\vec{\nabla} \cdot \vec{B} = 0, \quad \vec{\nabla} \cdot \vec{E} = 4\pi \rho_c, \quad (68)$$

$$\vec{\nabla} \times (\alpha_g \vec{E}) = - \left[\frac{\partial}{\partial t} - \mathcal{L}_{\vec{\beta}} \right] \vec{B}, \quad (69)$$

$$\vec{\nabla} \times (\alpha_g \vec{B}) = \left[\frac{\partial}{\partial t} - \mathcal{L}_{\vec{\beta}} \right] \vec{E} + 4\pi \alpha_g \vec{j}, \quad (70)$$

where $\mathcal{L}_{\vec{\beta}}$ is the Lie-derivative along $\vec{\beta}$, e.g. $\mathcal{L}_{\vec{\beta}} \vec{E} = (\vec{\beta} \cdot \vec{\nabla}) \vec{E} - (\vec{E} \cdot \vec{\nabla}) \vec{\beta}$.

3.5.1 Applicability of MHD and quasi-neutrality

I have mentioned criteria that limit the applicability of MHD: (i) Relativistically hot plasma and (ii) if the collision timescale of electrons becomes longer than dynamical timescales. The latter happens for low densities and high temperature, unless the collision rate is anomalous, or in a FIDO frame close to the horizon where $\gamma/\nu_c \rightarrow \infty$. As a muon gas that becomes 'decayless', if it moves with high Lorentz factor through an observer's frame, a plasma becomes collisionless in the frame of a FIDO close to the horizon. If case (ii) applies (e.g. if the Debye length becomes

very large) the assumption of quasi-neutrality should also break down.

4 CONCLUSION

The full set of MHD equations, including a generalized Ohm's law, for an inviscid fully ionized electron-ion plasma have been derived from a two-component plasma theory within the framework of the 3+1 split of the Kerr metric. It has been argued that the plasma can only be described as a single fluid, if its components are cold (i.e. specific internal energy density and pressure are negligible compared to the rest-mass energy density). This places constraints on the applicability of not only MHD but also of hydrodynamics to astrophysical objects such as hot relativistic jets and the hot regime of transsonic accretion flows onto black holes. Very close to the horizon FIDOs can no longer describe a plasma as a fluid, because the plasma particles become collisionless on local dynamical timescales.

The generalized Ohm's law is found to be equivalent to the special-relativistic expression, if the plasma is quasi-neutral in its rest frame. Gravitomagnetic effects of the order of 'current acceleration' terms appear explicitly only in charged plasma, or if the MHD-approximation of synchronized bulk and gravitomagnetic acceleration of the plasma components is not applied.

Unlike Ardavan (1976) and Blackman & Field (1993), who have derived their special-relativistic generalized Ohm's laws from invariant distribution functions, I have derived the generalized Ohm's law by combining the equations of motion for electrons and ions, thereby compromising details of the interaction between electrons and ions. The ansatz made for the collision term is, however, confirmed by yielding, relativistically correctly, the conduction current.

In a companion paper (Khanna 1998) I will apply this theory to describe a gravitomagnetic battery, which operates in plasma surrounding a Kerr black hole.

ACKNOWLEDGEMENTS

I gratefully acknowledge fruitful discussions with Max Camenzind, Harald Lesch, Jochen Peitz, Stefan Spindeldreher and Markus Thiele. This work is supported by the Deutsche Forschungsgemeinschaft (SFB 328).

REFERENCES

- Ardavan, H., 1976, ApJ 203, 226
 Bekenstein, J., Eichler, D., 1985, ApJ 298, 493
 Bekenstein, J., Oron, E., 1978, Phys. Rev. D 18, No. 6, 1809
 Beskin, V.S., Par'ev, V.I., 1993, Phys. Uspekhi 36 (6), 529
 Blackman, E.G., Field, G.B., 1993, Phys. Rev. Let. 71, No. 21, 3481
 Blandford, R.D., Znajek, R.L., 1977, MNRAS 179, 433
 Camenzind, M., 1986, A&A 162, 32
 Camenzind, M., 1987, A&A 184, 341
 Cowling T.G., 1934, MNRAS 94, 39
 Fendt, C., 1997, A&A, 319, 1025
 Khanna, R., 1997, in: Proceedings of the Second International Sakharov Conf. on Phys., Dremine, I.M., Semikhatov, A.M. (Eds.), World Scientific, p. 134

- Khanna, R., 1998, MNRAS, in press
 Khanna, R., Camenzind, M., 1994, ApJ 435, L129
 Khanna, R., Camenzind, M., 1996a, A&A 307, 665
 Khanna, R., Camenzind, M., 1996b, A&A 313, 1028
 Kippenhahn, R., Möllenhoff, C., 1975, Elementare Plasmaphysik, B.I.-Wissenschaftsverlag
 Kudoh, T., Kaburaki, O., 1996, ApJ 460, 199
 Okamoto, I., 1992, MNRAS 254, 192
 Phinney, E.S., Ph.D. diss., Univ. of Cambridge
 Schlüter, A., 1950, Z. Naturforschung 5a, 72
 Shapiro, S.L., Teukolsky, S.A., 1983, Black Holes, White Dwarfs and Neutron Stars, Wiley-Interscience, New York
 Spitzer, L., 1962, Physics of Fully Ionized Gases, Interscience, New York
 Takahashi, M., Nitta, S., Tatematsu, Y., Tomimatsu, A., 1990, ApJ 363, 206
 Thorne, K.S., Macdonald, D.A., 1982, MNRAS 198, 339
 Thorne, K.S., Price, R.H., Macdonald, D.A., Suen, W.-M., Zhang, X.-H., 1986, in: Black Holes: The Membrane Paradigm, Thorne, K.S., Price, R.H., Macdonald, D.A., (Eds.), Yale Univ. Press, New Haven

APPENDIX A: TRANSFORMATION OF CHARGE DENSITIES

Charge densities in the FIDO frame are given by $\rho_{cx} = -J_x^\mu U_\mu$. The transformation of charge densities between FIDO frame and some other frame follows from expressing the FIDO 4-velocity U_μ in terms of the other frame's 4-velocity $W_{o\mu}$ and its velocity as measured by FIDOs $v_{o\mu}$ (see Khanna & Camenzind 1996a). The transformation between FIDO frame and plasma rest frame K' is given by

$$\rho_{cx} = -J_x^\mu \left(\frac{W_\mu}{\gamma} - v_\mu \right) = \frac{\rho'_{cx}}{\gamma} + J_x^\mu v_\mu = \frac{\rho'_{cx}}{\gamma} + \vec{j}_x \cdot \vec{v}. \quad (A1)$$

The last equality follows by applying the definition of the current density measured by FIDOs $j_x^\mu = h^\mu_\nu J_x^\nu$:

$$J_x^\mu v_\mu = \rho_{cx} U^\mu v_\mu + j_x^\mu v_\mu = j_x^\mu v_\mu. \quad (A2)$$

Here the subscript 'x' may be 'i' for ions, 'e' for electrons or *blank* for the plasma. Inserting Eqs. (21) into Eq. (A1) yields the second equalities in Eqs. (20). The transformations between FIDO frame and electron rest frame K^e , or ion rest frame K^i , respectively, follow by inserting $U_\mu = W_{x\mu}/\gamma_x - v_{x\mu}$ into $\rho_{cx} = -J_x^\mu U_\mu$, where now $x=i,e$. This yields $\rho_{cx} = \rho_{cx}^x \gamma_x$.

APPENDIX B: PARTICLE CONSERVATION IN THE FIDO FRAME

Particle conservation is expressed in the spacetime viewpoint by $N_{x;\alpha}^\alpha = 0$, where the particle current $N_x^\alpha = n_x W_x^\alpha$ and W_x^α is the 4-velocity of the species 'x'. N_x^α splits into

$$n_x^\alpha = h^\alpha_\nu N_x^\nu = n_x \gamma_x v_x^\alpha \text{ and } \rho_{nx} = -U_\nu N_x^\nu = n_x \gamma_x. \quad (B1)$$

Therefore

$$N_{x;\alpha}^\alpha = (\rho_{nx} U^\alpha + n_x^\alpha)_{;\alpha} = 0, \quad (B2)$$

which leads, under observance of the relation for the determinants of 4-metric and 3-metric $g = -\alpha_g^2 h$ and multiplication with m_x , to Eq. (22).

APPENDIX C: SYNCHRONIZED BULK ACCELERATION OF IONS AND ELECTRONS

Adding Eqs. (36) (x=i,e) and using definition (40) yields on the l.h.s.

$$\begin{aligned} \rho_{mi}^i \frac{d(\gamma_i \vec{v}_i)}{d\tau_i} + \rho_{me}^e \frac{d(\gamma_e \vec{v}_e)}{d\tau_e} &= \rho_{mi}^i \frac{d(\gamma_i \vec{v}_i)}{d\tau_i} \\ &+ \rho_{me}^e \left(\frac{d(\gamma_i \vec{v}_i)}{d\tau_i} - \left(\frac{d(\gamma_i \vec{v}_i)}{d\tau_i} - \frac{d(\gamma_e \vec{v}_e)}{d\tau_e} \right) \right) \\ &\approx (\rho_{mi}^i + \rho_{me}^e) \frac{d(\gamma_i \vec{v}_i)}{d\tau_i} \approx \rho_m^i \frac{d(\gamma \vec{v})}{d\tau_p}, \end{aligned} \quad (C1)$$

which is the l.h.s. of Eq. (37). The following assumptions have been made in order to arrive at the last line: (i) Inequation (41) has been used and (ii) approximation (35) has been applied.

APPENDIX D: MANIPULATING

$$(Zm_e \gamma_e^2 \vec{j}_i - m_i \gamma_i^2 \vec{j}_e)$$

Equation (46) contains the term $(Zm_e \gamma_e^2 \vec{j}_i - m_i \gamma_i^2 \vec{j}_e) \times \vec{B}$. In Eq. (50) this term has mutated into $Zen_i m_i \gamma^2 (\gamma_i \vec{v} - \vec{j}/Zen_i) \times \vec{B}$. This happened in the following way: The equations

$$\vec{j} = \vec{j}_i + \vec{j}_e = \rho_{ci} \vec{v}_i + \rho_{ce} \vec{v}_e \quad (D1)$$

and

$$\gamma^2 \rho_m^i \vec{v} \approx \gamma_i^2 \rho_{mi}^i \vec{v}_i + \gamma_e^2 \rho_{me}^e \vec{v}_e \quad (D2)$$

can be combined to give

$$m_i \gamma_i^2 \vec{j}_e \approx m_i \gamma_i^2 \vec{j} - Ze \gamma_i \gamma^2 \rho_m^i \vec{v} \quad (D3)$$

and

$$Zm_e \gamma_e^2 \vec{j}_i \approx Z \rho_{ci} m_e \gamma_e^2 \vec{v} - \frac{Zm_e \rho_{me}^e \gamma_e^4 \rho_{ci}}{\rho_{mi}^i \gamma_i^2 \rho_{ce}} \vec{j}, \quad (D4)$$

where approximations of the order m_e/m_i and as given in Eq. (35) have been made. Putting things together and making the same kind of approximations yields the desired result.

APPENDIX E: GENERALIZED OHM'S LAW WITHOUT SYNCHRONIZED ACCELERATION

If the approximations of synchronized bulk and gravitomagnetic acceleration (Eqs. [41] and [42]) are not applied to Eq. (46), and if $\gamma_x \vec{v}_x$ in the bulk acceleration terms and $\vec{v}_x$ in the gravitomagnetic force term are eliminated with the aid of Eqs. (D3) and (D4), one gets instead of the l.h.s. of Eq. (47) (or instead of the last line of Eq. [50] with opposite sign)

$$\begin{aligned} \frac{Zn_i \gamma_e m_e}{n_e e \gamma_i^2} \left[\frac{\gamma_i}{\alpha_g} \frac{\partial}{\partial t} + \left(\gamma \vec{v} - \frac{\gamma_i}{\alpha_g} \vec{\beta} \right) \cdot \vec{\nabla} \right] (\gamma \vec{v}) \\ - \frac{m_e}{e \gamma_e} \left[\frac{\gamma_e}{\alpha_g} \frac{\partial}{\partial t} + \left(\frac{Zn_i}{n_e} \gamma \vec{v} - \frac{\vec{j}}{en_e} - \frac{\gamma_e}{\alpha_g} \vec{\beta} \right) \cdot \vec{\nabla} \right] \frac{Zn_i}{n_e} \gamma \vec{v} \\ + \frac{m_e}{e \gamma_e} \left[\frac{\gamma_e}{\alpha_g} \frac{\partial}{\partial t} + \left(\frac{Zn_i}{n_e} \gamma \vec{v} - \frac{\vec{j}}{en_e} - \frac{\gamma_e}{\alpha_g} \vec{\beta} \right) \cdot \vec{\nabla} \right] \frac{\vec{j}}{en_e} \\ - \frac{m_e}{e \gamma_e} \frac{\gamma_e}{\alpha_g} \vec{\nabla} \cdot \vec{\beta} \cdot \left(\frac{Zn_i}{n_e} (\gamma_e - \gamma_i) \vec{v} + \frac{\vec{j}}{en_e} \right). \end{aligned} \quad (E1)$$

One can recognize acceleration terms, current acceleration terms and gravitomagnetic terms as well. If the plasma is quasi-neutral and for $\gamma_e \approx \gamma_i \approx \gamma$, the first line and the second line (except $\vec{j} \cdot \vec{\nabla}(\gamma \vec{v})$) cancel.

It is important to note that, any extra term containing $\vec{j}$ or $\vec{B}$ in Ohm's law may allow for dynamo action. In particular, if such a term possessed an axisymmetric toroidal component that does not vanish in a neutral point of an axisymmetric poloidal magnetic field loop, it might support the growth of axisymmetric magnetic field. This would violate Cowling's theorem (Cowling 1934; Khanna & Camenzind 1996a).

The gravitomagnetic terms $(\vec{\beta} \cdot \vec{\nabla}) \vec{j}/en_e$ and $\vec{\nabla} \vec{\beta} \cdot \vec{j}/en_e$ do, in fact, individually possess such terms ($\pm \omega(\vec{j}_p \cdot \vec{\nabla} \ln \tilde{\omega}) \vec{e}_\phi/en_e$), but they cancel identically. If inequalities (41) (approximation of synchronized bulk acceleration) were applied to Eq. (46), without approximating the gravitomagnetic force terms in the same way (Eq. [42]), the only gravitomagnetic term in $\vec{j}$ that remained, would be the $\vec{\nabla} \vec{\beta} \cdot \vec{j}$ -term of Eq. (E1). This would allow for an axisymmetric gravitomagnetic dynamo, but would be an artefact of an incomplete approximation. The generalized Ohm's law does therefore not introduce a new axisymmetric *gravitomagnetic* dynamo.

The current acceleration terms are of the order

$$\frac{4\pi}{\omega_{pe}^2} \frac{d\vec{j}}{d\tau_p}. \quad (E2)$$

They will, in general, possess axisymmetric toroidal components, but compared to the conduction current $\vec{j}/\sigma\gamma_e$ they are of the order $(\nu_e \tau_p)^{-1}$, i.e. they violate Cowling's theorem only on timescales of the electron collision timescale, where the applicability of MHD breaks down.